Author Index

Alberts, G., Boomsma, F., Man in 't Veld, A. J. and Schalekamp, M. A. D. H.
Simultaneous determination of catecholamines and dobutamine in human plasma and urine by high-performance liquid chromatography with fluorimetric detection 583(1992)236

Alpert, A. J., see Boutin, J. A. 583(1992)137

Arrowood, S., Hoyt, Jr., A. M. and Sepaniak, M. J.
Monitoring of benzylpenicillin decomposition in gastric contents by capillary zone electrophoresis 583(1992)105

Beijnen, J. H., see Kaijser, G. P. 583(1992)175

Bellon, B., see Rugani, N. 583(1992)246

Benoit, E., Jaussaud, P., Besse, S., Videmann, B., Courtot, D., Delatour, P. and Bonnaire, Y.
Identification of a benzhydrolic metabolite of ketoprofen in horses by gas chromatography–mass spectrometry and high-performance liquid chromatography 583(1992)167

Benowitz, N. L., see Jacob, III, P. 583(1992)145

Berlin, K. D., see Chen, C. L. 583(1992)274

Besse, S., see Benoit, E. 583(1992)167

Bonnaire, Y., see Benoit, E. 583(1992)167

Boomsma, F., see Alberts, G. 583(1992)236

Boutin, J. A., Ernould, A.-P., Ferry, G., Genton, A. and Alpert, A. J.
Use of hydrophilic interaction chromatography for the study of tyrosine protein kinase specificity 583(1992)137

Brocks, D. R., see Wright, M. R. 583(1992)259

Brune, K., see Schmidt, N. 583(1992)195

Bult, A., see Kaijser, G. P. 583(1992)175

Callebat, I., see Maisonneuve, A. 583(1992)155

Chen, C. L., Lesseley, B. A., Clarke, C. R., Roder, J. D., Sangiah, S., Berlin, K. D., Garrison, G. L., Scherlag, B. J., Lazzara, R. and Patterson, E.
High-performance liquid chromatographic determination of BRB-I-28, a novel antiarrhythmic agent, in dog plasma and urine 583(1992)274

Chmelík, J., see Urbánková, E. 583(1992)27

Clarke, C. R., see Chen, C. L. 583(1992)274

Coppet, L., see Maisonneuve, A. 583(1992)155

Courtot, D., see Benoit, E. 583(1992)167

Cozzone, P. J., see Rugani, N. 583(1992)246

Debordes, L., see Maisonneuve, A. 583(1992)155

De Kraker, J., see Kaijser, G. P. 583(1992)175

De La Fournière, L., see Rugani, N. 583(1992)246

Delatour, P., see Benoit, E. 583(1992)167

Dezan, C., see Rugani, N. 583(1992)246

Diquet, B., see Thomare, P. 583(1992)217

Ducros, V. and Favier, A.
Gas chromatographic–mass spectrometric method for the determination of selenium in biological samples 583(1992)35

Edwards, G., see Thomas, C. G. 583(1992)131

Ernould, A.-P., see Boutin, J. A. 583(1992)137

Favier, A., see Ducros, V. 583(1992)35

Ferry, G., see Boutin, J. A. 583(1992)137

Flynn, J. W. and O'Brien, J. E.
High-performance liquid chromatographic method for the simultaneous measurement of remacemide (a novel anticonvulsant and cerebroprotectant) and an active metabolite in human plasma 583(1992)91

Fujimoto, K., see Okamoto, J. 583(1992)53

Garrison, G. L., see Chen, C. L. 583(1992)274

Geisslinger, G., see Schmidt, N. 583(1992)195

Genton, A., see Boutin, J. A. 583(1992)137

Gumbhir, K. and Mason, W. D.
High-performance liquid chromatographic method for the determination of tris(hydroxymethyl)aminomethane (tromethamine) in human plasma 583(1992)99

Haagsma, N., see Moretti, V. M. 583(1992)77

Hadjimichael, J., see Rossomando, E. F. 583(1992)11

Hadjimichael, J., see Rossomando, E. F. 583(1992)19

Hatada, K., Kimura, M., Ono, I. and Ozaki, M.
Determination of a new calcium antagonist and its main metabolite in plasma by thermospray liquid chromatography–mass spectrometry 583(1992)116

Heykants, J., see Woestenborghs, R. 583(1992)223

Hiemke, C., see Weigmann, H. 583(1992)209

Hirai, E., see Okamoto, J. 583(1992)45

Hirai, E., see Okamoto, J. 583(1992)53

Hormazabal, V. and Rogstad, A.
Simultaneous determination of sulphadiazine and trimethoprim in plasma and tissues of cultured fish for residual and pharmacokinetic studies 583(1992)201

Hoyt, Jr., A. M., see Arrowood, S. 583(1992)105

Huupponen, R., Ojala-Karlsson, P., Rouru, J. and Koulu, M.
Determination of metformin in plasma by high-performance liquid chromatography 583(1992)270

Jacob, III, P., Shulgin, A. T., Yu, L. and Benowitz, N. L.
Determination of the nicotine metabolite *trans*-3′-hydroxycotinine in urine of smokers using gas chromatography with nitrogen-selective detection or selected ion monitoring 583(1992)145

Jamali, F., see Wright, M. R. 583(1992)259

Jaussaud, P., see Benoit, E. 583(1992)167

Kaijser, G. P., Beijnen, J. H., Bult, A., Wiese, G., De Kraker, J., Keizer, H. J. and Underberg, W. J. M.
Gas chromatographic determination of 2- and 3-dechloroethylifosfamide in plasma and urine 583(1992)175

Kajbaf, M., Sepai, O., Lamb, J. H. and Naylor, S.
Identification of metabolites of 4,4′-diaminodiphenylmethane (methylene dianiline) using liquid chromatographic and mass spectrometric techniques 583(1992)63

Kakinuma, H., see Yokoyama, Y. 583(1992)1

Keizer, H. J., see Kaijser, G. P. 583(1992)175

Kikuta, C., see Mascher, H. 583(1992)122

Kimura, M., see Hatada, K. 583(1992)116

Kitagawa, T., see Okamoto, J. 583(1992)45

Kitagawa, T., see Okamoto, J. 583(1992)53

Kline, W. F. and Matuszewski, B. K.
Improved determination of the bisphosphonate alendronate in human plasma and urine by automated precolumn derivatization and high-performance liquid chromatography with fluorescence and electrochemical detection 583(1992)183
Koulu, M., see Huupponen, R. 583(1992)270
Kumashiro, S., see Watanabe, A. 583(1992)241
Kurumaji, A., see Watanabe, A. 583(1992)241
Lamb, J. H., see Kajbaf, M. 583(1992)63
Lazzara, R., see Chen, C. L. 583(1992)274
Lennard, L. and Singleton, H. J.
High-performance liquid chromatographic assay of the methyl and nucleotide metabolites of 6-mercaptopurine: quantitation of red blood cell 6-thioguanine nucleotide, 6-thioinosinic acid and 6-methylmercaptopurine metabolites in a single sample 583(1992)83
Lesseley, B. A., see Chen, C. L. 583(1992)274
Lorreyne, W., see Woestenborghs, R. 583(1992)223
Maisonneuve, A., Callebat, I., Debordes, L. and Coppet, L.
Specific and sensitive quantitation of 2,2′-dichlorodiethyl sulphide (sulphur mustard) in water, plasma and blood: application to toxicokinetic study in the rat after intravenous intoxication 583(1992)155
Man in 't Veld, A. J., see Alberts, G. 583(1992)236
Mascher, H., Kikuta, C., Metz, R. and Vergin, H.
New, high-sensitivity high-performance liquid chromatographic method for the determination of acyclovir in human plasma, using fluorometric detection 583(1992)122
Mason, W. D., see Gumbhir, K. 583(1992)99
Matsubara, T., see Okamoto, J. 583(1992)45
Matsubara, T., see Okamoto, J. 583(1992)53
Matulík, F., see Urbánková, E. 583(1992)27
Matuszewski, B. K., see Kline, W. F. 583(1992)183
McCallister, J. D., see Patrick, K. S. 583(1992)254
Metz, R., see Mascher, H. 583(1992)122
Moncrieff, J.
Determination of cetirizine in serum using reversed-phase high-performance liquid chromatography with ultraviolet spectrophotometric detection 583(1992)128
Moretti, V. M., Van de Water, C. and Haagsma, N.
Automated high-performance liquid chromatographic determination of chloramphenicol in milk and swine muscle tissue using on-line immunoaffinity sample clean-up 583(1992)77
Nakano, S., see Noguchi, H. 583(1992)266
Naruto, S., see Noguchi, H. 583(1992)266
Naylor, S., see Kajbaf, M. 583(1992)63
Nguyen, B. L., see Patrick, K. S. 583(1992)254
Noguchi, H., Tomita, N., Naruto, S. and Nakano, S.
Determination of gliclazide in serum by high-performance liquid chromatography using solid-phase extraction 583(1992)266
Nováková, N., see Urbánková, E. 583(1992)27
O'Brien, J. E., see Flynn, J. W. 583(1992)91
Ojala-Karlsson, P., see Huupponen, R. 583(1992)270
Okamoto, J., Hirai, E., Kitagawa, T. and Matsubara, T.
Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human plasma 583(1992)45
Okamoto, J., Hirai, E., Suzuki, Y., Fujimoto, K., Kitagawa, T. and Matsubara, T.
Capillary gas chromatographic method for the determination of the thromboxane A_2 receptor antagonist S-1452 and its metabolites in human urine 583(1992)53
Oliw, E. H.
Enantioselective separation of some polyunsaturated epoxy fatty acids by high-performance liquid chromatography on a cellulose phenylcarbamate (Chiralcel OC) stationary phase 583(1992)231
Ono, I., see Hatada, K. 583(1992)116
Ozaki, M., see Hatada, K. 583(1992)116
Patrick, K. S., Nguyen, B. L. and McCallister, J. D.
Gas chromatographic–mass spectrometric determination of plasma selegiline using a deuterated internal standard 583(1992)254
Patterson, E., see Chen, C. L. 583(1992)274
Purdy, W. C., see Tadey, T. 583(1992)111
Roder, J. D., see Chen, C. L. 583(1992)274
Rogstad, A., see Hormazabal, V. 583(1992)201
Rossomando, E. F., White, L. B., Hadjimichael, J. and Shafer, D.
Immunomagnetic separation of tumor necrosis factor α. I. Batch procedure for human temporomandibular fluid 583(1992)11
Rossomando, E. F., White, L. B. and Hadjimichael, J.
Immunomagnetic separation of tumor necrosis factor α. II. *In situ* procedure for the human gingival space 583(1992)19
Rouru, J., see Huupponen, R. 583(1992)270
Rugani, N., Dezan, C., De La Fournière, L., Cozzone, P. J., Bellon, B. and Sarda, L.
Separation and characterization of the precursor and activated forms of porcine and human pancreatic colipase by reversed-phase liquid chromatography 583(1992)246
Sangiah, S., see Chen, C. L. 583(1992)274
Sarda, L., see Rugani, N. 583(1992)246
Sato, H., see Yokoyama, Y. 583(1992)1
Sattari, S., see Wright, M. R. 583(1992)259
Schalekamp, M. A. D. H., see Alberts, G. 583(1992)236
Scherlag, B. J., see Chen, C. L. 583(1992)274
Schmidt, N., Brune, K. and Geisslinger, G.
Stereoselective determination of the enantiomers of methadone in plasma using high-performance liquid chromatography 583(1992)195
Semba, J., see Watanabe, A. 583(1992)241
Sepai, O., see Kajbaf, M. 583(1992)63
Sepaniak, M. J., see Arrowood, S. 583(1992)105
Shafer, D., see Rossomando, E. F. 583(1992)11
Shulgin, A. T., see Jacob, III, P. 583(1992)145
Singleton, H. J., see Lennard, L. 583(1992)83
Suzuki, Y., see Okamoto, J. 583(1992)53
Tadey, T. and Purdy, W. C.
Characterization of plasma apolipoproteins by capillary electrophoresis 583(1992)111
Thomare, P., Wang, K., Van Der Meersch-Mougeot, V. and Diquet, B.
Sensitive micromethod for column liquid chromatographic determination of fluoxetine and norfluoxetine in human plasma 583(1992)217

Thomas, C. G., Ward, S. A. and Edwards, G.
Selective determination, in plasma, of artemether and its major metabolite, dihydroartemisinin, by high-performance liquid chromatography with ultraviolet detection 583(1992)131
Tomita, N., see Noguchi, H. 583(1992)266
Toru, M., see Watanabe, A. 583(1992)241
Tsuchiya, M., see Yokoyama, Y. 583(1992)1
Underberg, W. J. M., see Kaijser, G. P. 583(1992)175
Urbánková, E., Vacek, A., Nováková, N., Matulík, F. and Chmelík, J.
Investigation of red blood cell fractionation by gravitational field-flow fractionation 583(1992)27
Vacek, A., see Urbánková, E. 583(1992)27
Van Der Meersch-Mougeot, V., see Thomare, P. 583(1992)217
Van de Water, C., see Moretti, V. M. 583(1992)77
Van Rompaey, F., see Woestenborghs, R. 583(1992)223
Vergin, H., see Mascher, H. 583(1992)122
Videmann, B., see Benoit, E. 583(1992)167
Wang, K., see Thomare, P. 583(1992)217
Ward, S. A., see Thomas, C. G. 583(1992)131
Watanabe, A., Semba, J., Kurumaji, A., Kumashiro, S. and Toru, M.
Measurement of glutamate, aspartate and glycine and its potential precursors in human brain using high-performance liquid chromatography by pre-column derivatization with diethylaminoazobenzene sulphonyl chloride 583(1992)241
Weigmann, H. and Hiemke, C.
Determination of clozapine and its major metabolites in human serum using automated solid-phase extraction and subsequent isocratic high-performance liquid chromatography with ultraviolet detection 583(1992)209
White, L. B., see Rossomando, E. F. 583(1992)11
White, L. B., see Rossomando, E. F. 583(1992)19
Wiese, G., see Kaijser, G. P. 583(1992)175
Woestenborghs, R., Lorreyne, W., Van Rompaey, F. and Heykants, J.
Determination of risperidone and 9-hydroxyrisperidone in plasma, urine and animal tissues by high-performance liquid chromatography 583(1992)223
Wright, M. R., Sattari, S., Brocks, D. R. and Jamali, F.
Improved high-performance liquid chromatographic assay method for the enantiomers of ibuprofen 583(1992)259
Yokoyama, Y., Tsuchiya, M., Sato, H. and Kakinuma, H.
Determination of creatinine and ultraviolet-absorbing amino acids and organic acids in urine by reversed-phase high-performance liquid chromatography 583(1992)1
Yu, L., see Jacob, III, P. 583(1992)145